arXiv:hep-th/9903220v1 25 Mar 1999

ON THE BRACKET PROBLEM IN COVARIANT HAMILTONIAN FIELD THEORY

L.Mangiarotti
Department of Mathematics and Physics, Camerino University, 62032 Camerino, Italy
E-mail: mangiaro@camserv.unicam.it

G.Sardanashvily
Department of Theoretical Physics, Moscow State University
117234 Moscow, Russia
E-mail: sard@grav.phys.msu.su

Abstract

The polysymplectic phase space of covariant Hamiltonian field theory can be provided with the current algebra bracket.

As is well known, when applied to field theory, the familiar symplectic technique of mechanics takes the form of instantaneous Hamiltonian formalism on an infinite-dimensional phase space. The finite-dimensional covariant Hamiltonian approach to field theory is vigorously developed from the seventies in its multisymplectic and polysymplectic variants (see [4, 5, 6, 13] for a survey). They are related to the two different Legendre morphisms in the first order calculus of variations on fiber bundles.

Recall that, given a fiber bundle $Y \to X$, coordinated by (x^λ, y^i), a first order Lagrangian L is defined as a semibasic density

$$L = \mathcal{L}\omega : J^1Y \to \overset{n}{\wedge} T^*X, \quad \omega = dx^1 \wedge \cdots dx^n, \quad n = \dim X,$$

on the affine jet bundle $J^1Y \to Y$, provided with the adapted coordinates $(x^\lambda, y^i, y^i_\lambda)$. J^1Y can be seen as a finite-dimensional configuration space of fields represented by sections of $Y \to X$. The Poincaré–Cartan form

$$H_L = \mathcal{L}\omega + \pi^\lambda_i(dy^i - y^i_\mu dx^\mu) \wedge \omega_\lambda, \quad \pi^\lambda_i = \partial^\lambda_i \mathcal{L}, \quad \omega_\lambda = \partial_\lambda \rfloor \omega,$$

can be defined as a Lepagean equivalent of L which is a semibasic form on $J^1Y \to Y$. It yields the Legendre bundle morphism $\widehat{H}_L$ over Y of J^1Y to the homogeneous Legendre bundle

$$Z_Y = J^{1\star}Y = T^*Y \wedge (\overset{n-1}{\wedge} T^*X) \tag{1}$$

which is the affine $\overset{n+1}{\wedge}$-valued dual of $J^1Y \to Y$, and is endowed with the holonomic coordinates $(x^\lambda, y^i, p_i^\lambda, p)$. In these coordinates, the morphism (1) reads

$$(p_i^\mu, p) \circ \widehat{H}_L = (\pi_i^\mu, \mathcal{L} - \pi_i^\mu y_\mu^i).$$

The fiber bundle Z_Y is treated as a homogeneous finite-dimensional phase space of fields. It is equipped with the canonical multisymplectic form

$$d\Xi_Y = dp \wedge \omega + dp_i^\lambda \wedge dy^i \wedge \omega_\lambda.$$

Every Lagrangian L defines the Legendre map $\widehat{L}$ over Y of J^1Y to the Legendre bundle

$$\Pi = \overset{n}{\wedge} T^*X \underset{Y}{\otimes} V^*Y \underset{Y}{\otimes} TX \tag{2}$$

provided with the holonomic coordinates $(x^\lambda, y^i, p_i^\lambda)$. Π plays the role of a finite-dimensional momentum phase space of fields. It is equipped with the canonical polysymplectic form

$$\Omega_Y = dp_i^\lambda \wedge dy^i \wedge \omega \otimes \partial_\lambda.$$

The relationship between the multisymplectic and polysymplectic phase spaces is given by the exact sequence

$$0 \to \Pi \underset{X}{\times} \overset{n}{\wedge} T^*X \hookrightarrow Z_Y \to \Pi \to 0,$$

where $\pi_{Z\Pi} : Z_Y \to \Pi$ is a 1-dimensional affine bundle. Given a section h of $Z_Y \to \Pi$, the pull-back $h^*\Xi_Y$ is a polysymplectic Hamiltonian form on Π [3, 4, 13].

A natural idea is to generalize a Poisson bracket in symplectic mechanics to multisymplectic or polysymplectic manifolds and, as a final result, to come to the covariant canonical quantization of field theory. Different variants of such a bracket have been suggested (see [9, 10] and references therein). The main difficulty is that the bracket must be globally defined. Let us note that multisymplectic manifolds, whose particular example is $(Z_Y, d\Xi_Y)$, look rather promising for algebraic constructions since multisymplectic forms are exterior forms [1, 2, 7, 8, 11].

Nevertheless, the $X = \mathbf{R}$ reduction of the covariant Hamiltonian formalism leads to time-dependent mechanics, but not conservative symplectic mechanics [12, 15]. In this case, $Z_Y = T^*Y$ and $\Pi = V^*Y$ are homogeneous and momentum phase spaces of time-dependent mechanics, respectively. The momentum phase space V^*Y, coordinated by $(t, y^i, p_i = \dot{y}_i)$, is endowed with the canonical degenerate Poisson structure given by the Poisson bracket

$$\{f, g\}_V = \partial^i f \partial_i g - \partial^i g \partial_i f, \qquad f, g \in C^\infty(V^*Y). \tag{3}$$

However, a Poisson bracket $\{\mathcal{H}, f\}_V$ of a Hamiltonian $\mathcal{H}$ and functions f on the momentum phase space V^*Y fails to be a well-behaved entity because a Hamiltonian of time-dependent mechanics is not a scalar with respect to time-dependent transformations, e.g., time-dependent canonical transformations. In particular, the equality $\{\mathcal{H}, f\}_V = 0$ is not preserved under time-dependent transformations. As a consequence, the evolution equation in time-dependent mechanics is not reduced to a Poisson bracket, and integrals of motion are not functions in involution with a Hamiltonian [12, 15].

At the same time, the Poisson bracket (3) leads to the following current algebra bracket. Let $u = u^i \partial_i$ be a vertical vector field on $Y \to \mathbf{R}$, and $J_u = u^i p_i$ the corresponding symmetry current on V^*Y along u. In particular, J_u is a conserved Nöther current if a Hamiltonian form

$$H = p_i dy^i - \mathcal{H} dt$$

is invariant under the 1-parameter group of local gauge transformations whose generator is the vector field u [12, 15]. The symmetry currents J_u constitute a Lie algebra with respect to the bracket

$$[J_u, J_{u'}] = \{J_u, J_{u'}\}_V = J_{[u,u']}.$$

This current algebra bracket can be extended to the general polysymplectic case as follows.

There is the canonical isomorphism

$$\overline{\theta} = p_i^\lambda \overline{d} y^i \wedge \omega_\lambda : \Pi \to V^*Y \underset{Y}{\wedge} (\overset{n-1}{\wedge} T^*X),$$

where $\{\overline{d}y^i\}$ are the fiber bases for the vertical cotangent bundle V^*Y of $Y \to X$. Let $u = u^i \partial_i$ be a vertical vector field on $Y \to X$. The corresponding symmetry current is defined as a semibasic exterior $(n-1)$-form

$$J_u = u\rfloor\overline{\theta} = u^i p^\lambda_i \omega_\lambda \tag{4}$$

on the Legendre bundle Π (2). In particular, J_u is a Nöther current if a Hamiltonian form H is invariant under the 1-parameter group of local gauge transformations whose generator is the lift onto Π of the vector field u [4, 14]. The symmetry currents (4) constitute a Lie algebra with respect to the bracket

$$[J_u, J_{u'}] \overset{\text{def}}{=} J_{[u,u']}. \tag{5}$$

If $Y \to X$ is a vector bundle and X is provided with a non-degenerate metric g, the bracket (5) can be extended to any semibasic exterior $(n-1)$-forms $\phi = \phi^\alpha\omega_\alpha$ on Π by the law

$$[\phi,\sigma] = g_{\alpha\beta}g^{\mu\nu}(\partial^i_\mu\phi^\alpha\partial_i\sigma^\beta - \partial^i_\mu\sigma^\beta\partial_i\phi^\alpha)\omega_\nu.$$

Similarly, the bracket of semibasic 1-forms on Π is defined [12].

The bracket (5) looks promising for the current algebra quantization of the covariant Hamiltonian formalism.

References

1. Cantrijn, F., Ibort, L.A., and de León, M.: Hamiltonian structures on multisymplectic manifolds. Rend. Sem. Mat. Univ. Pol. Torino. **54**, 225 (1996)

2. Cantrijn, F., Ibort, A. and de León, M. On the geometry of multisymplectic manifolds. J. Australian Math. Soc. A (to appear)

3. Cariñena, J., Crampin M. and Ibort, L.A.: On the multisymplectic formalism for first order field theories. Diff. Geom. and Appl. **1**, 345 (1991)

4. Giachetta, G., Mangiarotti, L. and Sardanashvily, G.: New Lagrangian and Hamiltonian methods in field theory. Singapore: World Scientific 1997

5. Gotay, M. A multisymplectic framework for classical field theory and the calculus of variations. I. Covariant Hamiltonian formalism. In: M.Francaviglia (ed.) Mechanics, analysis and geometry: 200 years after Lagrange. pp.203-235. Amsterdam: North-Holland 1991

6. Gotay, M., Isenberg, J. and Marsden, J.: Momentum maps and classical relativistic fields. E-print physics/9801019

7. Hrabak, S.: On a multisymplectic formulation of the classical BRST symmetry for first order field theories. E-prints math-ph/9901012;013

8. Ibort, A., Echeverría-Enríquez, A., Muñoz-Lecanda, M. and Román-Roy, N.: Invariant forms and automorphisms of multisymplectic manifolds. E-print math.DG/9805040

9. Kanatchikov, I.: Toward the Born-Weyl quantization of fields. Int. J. Theor. Phys. **37**, 333 (1998)

10. Kanatchikov, I.: On quantization of field theories in polymomentum variables. E-print hep-th/9811016

11. Lawson, J.: A frame bundle generalization of multisymplectic field theories. E-print dg-ga/9706008

12. Mangiarotti, L. and Sardanashvily, G.: Gauge mechanics. Singapore: World Scientific 1998

13. Sardanashvily, G.: Generalized Hamiltonian formalism for field theory. Constraint systems. Singapore: World Scientific 1995

14. Sardanashvily, G.: Stress-energy-momentum tensors in constraint field theories. J. Math. Phys. **38**, 847 (1997)

15. Sardanashvily, G.: Hamiltonian time-dependent mechanics. J. Math. Phys. **39**, 2714 (1998)